



CASE REPORT

# Case Report: Rare presentation of spinal meningioma in

## Kosovo

[version 1; peer review: awaiting peer review]

Flamur Hasimi<sup>1</sup>, Fjolla Ymeri<sup>2</sup><sup>1</sup>Neurosurgery Clinic, University Clinical Centre of Kosovo, Prishtina, Prishtina, Kosovo (Serbia and Montenegro)<sup>2</sup>Kolegji Universitar Universum, Prishtina, Prishtina, Kosovo (Serbia and Montenegro)

**V1** First published: 05 Jan 2026, 6:7  
<https://doi.org/10.12688/openreseurope.22197.1>

Latest published: 05 Jan 2026, 6:7  
<https://doi.org/10.12688/openreseurope.22197.1>

### Abstract

This case report outlines the diagnosis and management of a spinal meningioma in an 18-year-old female presenting with progressive weakness in both lower limbs. Magnetic resonance imaging revealed an intradural extramedullary tumour compressing the spinal cord, subsequently confirmed as a meningioma. Urgent neurosurgical intervention was performed, involving total laminectomy at the TH6/7 vertebral level and meticulous tumour resection. Postoperatively, the patient demonstrated significant neurological improvement, with muscle strength increasing from 2/5 to 4/5. This case highlights the importance of prompt diagnosis, multidisciplinary management, and close postoperative monitoring in achieving favourable outcomes for spinal meningiomas.

### Keywords

spinal meningioma, neurosurgical intervention, intradural extramedullary tumor, magnetic resonance imaging

### Open Peer Review

**Approval Status** *AWAITING PEER REVIEW*

Any reports and responses or comments on the article can be found at the end of the article.



This article is included in the [COST Actions gateway](#).

**Corresponding author:** Flamur Hasimi ([hasimiflamur0@gmail.com](mailto:hasimiflamur0@gmail.com))

**Author roles:** **Hasimi F:** Conceptualization, Formal Analysis, Funding Acquisition, Investigation, Methodology, Resources, Supervision, Validation, Visualization; **Ymeri F:** Software, Writing – Original Draft Preparation, Writing – Review & Editing

**Competing interests:** No competing interests were disclosed.

**Grant information:** This project has received funding from the European Union's Framework Programme for Research & Innovation as part of the COST Action CA22103 A COMPREHENSIVE NETWORK AGAINST BRAIN CANCER, as supported by the COST Association (European Cooperation in Science and Technology).

*The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.*

**Copyright:** © 2026 Hasimi F and Ymeri F. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

**How to cite this article:** Hasimi F and Ymeri F. **Case Report: Rare presentation of spinal meningioma in Kosovo [version 1; peer review: awaiting peer review]** Open Research Europe 2026, 6:7 <https://doi.org/10.12688/openreseurope.22197.1>

**First published:** 05 Jan 2026, 6:7 <https://doi.org/10.12688/openreseurope.22197.1>

## Introduction


Spinal meningiomas (SM) are growths that generally have a positive outlook both in terms of cancer treatment and surgical outcomes, with a rare chance of the tumour returning. Usually found in the space surrounding the spinal cord but outside its innermost layer, these tumours develop gradually and tend to extend sideways into the area around the spinal cord, often stretching the nearby arachnoid membrane but seldom affecting the innermost pia mater (Hohenberger *et al.*, 2023). Meningiomas are lesions that arise from arachnoidal cap cells, the outer part of the arachnoid layer and villi. They usually form dural attachments and are marked by meningothelial hyperplasia. Spinal meningiomas most commonly occur in the posterior, posterolateral, or lateral thoracic region, followed by the anterior cervical and lumbosacral regions (Sun *et al.*, 2020). The standard approach to treating them involves surgery, aiming to completely remove the tumour and enhance neurological function. The main treatment is surgery, which can be performed in (a) a classical open microsurgical approach in the vast majority of cases, (b) a minimally invasive surgery (MIS), or (c) through an endoscopic intervention (Serratrice *et al.*, 2023; Sun *et al.*, 2020).

## Case presentation

The patient, an 18-year-old female, with no relevant past medical history and no known family history of neurological or oncological disease, presented with complaints of progressive weakness in both lower limbs, with a recorded muscle strength of 2/5. She reported significant difficulty in walking and relied on a wheelchair for mobility. Upon examination, neurological deficits were noted, prompting further investigation. Written informed consent was also obtained from the patient for publication of this case report and accompanying images. All reasonable efforts were made to anonymize patient information.

## Diagnostic assessment

Diagnostic workup included magnetic resonance imaging (MRI) of the spinal axis, revealing an intradural extramedullary tumour compressing the spinal cord (Figure 1). The tumour



**Figure 1.** The image shows the magnetic resonance imaging (MRI) scan of the spinal axis, specifically highlighting an expansile intradural extramedullary process.

was identified as a meningioma, characterized by its firm consistency and white-gray appearance upon exposure during surgery. The preoperative MRI provided crucial insights into the location and extent of the tumour, guiding the surgical approach. Figure 1 shows a magnetic resonance imaging (MRI) scan of the spinal axis, specifically highlighting an expansile intradural extramedullary process.

## Treatment


The patient underwent urgent neurosurgical intervention to address the spinal meningioma. The surgical procedure involved total laminectomy at the TH6/7 vertebral level (Figure 2) to access and remove the tumour. Intraoperative neurosurgical procedure for the resection of a spinal meningioma is shown in Figure 2. Intraoperatively, meticulous care was taken to preserve neural structures while achieving complete tumour resection as shown in Figure 3. The excised mass was sent for histopathological examination to confirm the diagnosis and assess for any concerning features.

## Follow-up and outcome

Following surgery, the patient was closely monitored in the postoperative period for any signs of neurological improvement or complications. Gradually, the patient exhibited notable improvement in muscle strength, with postoperative muscle strength reaching 4/5 compared to the preoperative level of 2/5. Regular follow-up appointments were scheduled to monitor the patient's progress, ensuring optimal recovery and functional outcomes.

## Discussion

Although meningiomas are generally benign tumours, atypical and rare clinical presentations have been reported in the literature, emphasizing the need for individualized diagnostic and therapeutic approaches (Gone *et al.*, 2024). The successful management of this case underscores the efficacy of prompt neurosurgical intervention coupled with comprehensive preoperative imaging and meticulous tumour resection in achieving favourable outcomes for spinal meningiomas. The utilization of magnetic resonance imaging (MRI) facilitated accurate preoperative planning, enabling the surgical team to approach the tumour with precision while minimizing damage to



**Figure 2.** The image shows Intraoperative neurosurgical procedure for the resection of a spinal meningioma.



**Figure 3. Complete tumour resection.**

adjacent neural structures (Hohenberger *et al.*, 2023; Serratrice *et al.*, 2023). Total laminectomy at the TH6/7 vertebral level provided adequate exposure for complete tumour resection, with intraoperative vigilance ensuring preservation of neurological function. Previous studies have demonstrated that gross total resection is associated with lower recurrence rates and improved long-term neurological outcomes compared with subtotal resection (Materi *et al.*, 2020). The observed postoperative improvement in muscle strength highlights the potential for significant neurological recovery following surgical intervention for spinal meningiomas. Thoracic tumours often present with progressive neurological deficits, and early surgical management can lead to significant functional recovery (Sun *et al.*, 2020). Emerging evidence suggests that genomic subgroups of meningiomas may correlate with distinct clinical behaviours and outcomes, highlighting the potential role of molecular

profiling in future risk stratification and management strategies (Youngblood *et al.*, 2019). Regular follow-up appointments are essential for monitoring long-term outcomes and addressing any potential complications, emphasizing the importance of a multidisciplinary approach in optimizing patient care.

### Conclusion

This case exemplifies the successful management of a spinal meningioma through prompt neurosurgical intervention guided by preoperative imaging. The observed neurological improvement underscores the importance of surgical resection in restoring function in affected patients. Long-term follow-up is crucial for monitoring outcomes and addressing any potential complications. Continued research efforts are needed to further elucidate optimal treatment strategies and improve outcomes for patients with spinal meningiomas.

### Ethics and consent

Ethical approval not required. Written informed consent was obtained from the patient for participation in the study and for publication of this case report.

### Data availability

All data underlying the findings of this case report are included within the article. Additional clinical data are archived at the University Hospital Centre and are not publicly available due to ethical and privacy considerations, as they contain potentially identifiable patient information. The CARE checklist for this case report is available in Zenodo under the title “CARE checklist for Rare Presentation of Spinal Meningioma in Kosova: A Case Report”, DOI: [10.5281/zenodo.17969909](https://doi.org/10.5281/zenodo.17969909) licensed under CC0 (Hasimi & Ymeri, 2025).

### References

- Gone J, Kennedy B, Fontaine T, *et al.*: **Creeping through the cranium: a rare and interesting presentation of a meningioma.** *Radiol Case Rep.* 2024; **19**(9): 3928–3933.  
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Hasimi F, Ymeri F: **Rare presentation of spinal meningioma in Kosova: case report.** Zenodo. 2025.  
<http://www.doi.org/10.5281/ZENODO.17969909>
- Hohenberger C, Hau P, Schebesch KM, *et al.*: **Spinal meningiomas.** *Neurooncol Adv.* 2023; **5**(Suppl 1): i112–i121.  
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Materi J, Mampre D, Ehresman J, *et al.*: **Predictors of recurrence and high growth rate of residual meningiomas after subtotal resection.** *J Neurosurg.* 2020; **134**(2): 410–416.  
[PubMed Abstract](#) | [Publisher Full Text](#)
- Serratrice N, Lameche I, Attieh C, *et al.*: **Spinal meningiomas, from biology to management — A literature review.** *Front Oncol.* 2023; **12**: 1084404.  
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Sun C, Dou Z, Wu J, *et al.*: **The preferred locations of meningioma according to different biological characteristics based on voxel-wise analysis.** *Front Oncol.* 2020; **10**: 1412.  
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Youngblood MW, Duran D, Montejo JD, *et al.*: **Correlations between genomic subgroup and clinical features in a cohort of more than 3000 meningiomas.** *J Neurosurg.* 2019; **133**(5): 1345–1354.  
[PubMed Abstract](#) | [Publisher Full Text](#)